

Comment on: "Current-voltage characteristics and zero-resistance state in 2DEG"

M. V. Cheremisin

A.F.Ioffe Physical-Technical Institute, St.Petersburg, Russia

We demonstrate that N(S)-shape current-voltage characteristics proposed to explain zero-resistance state in Corbino(Hall bar) geometry 2DEG (cond-mat//0302063, 0303530) cannot account essential features of radiation-induced magnetoresistance oscillations experiments.

Recently, a great deal of interest has been focussed on the anomalous transport behavior of 2DEG subjected to microwave and weak magnetic fields [1, 2]. Possible scenarios for this phenomenon were discussed in [3]-[9]. As it was shown in [4, 5, 8], at certain magnetic field and irradiation strength the longitudinal conductivity may become negative argued to result in zero resistance(conductance) states ZRS(ZCS)observed in experiments. In Ref.[10] current-voltage characteristic (CVC) of irradiated 2D system in classically strong magnetic field suggested to exhibit N(S)-shaped behavior(Fig.1) in Corbino(Hall bar) geometry respectively.

We now discuss the relevance of the ZRS model[10] regarding to experiments [2, 11]. According to [5, 10], in ZRS the Hall bar sample exhibits the current domain structure(bold line in Fig.1,bottom panel), when $|j| < j_0$. Upon current enhancement $|j| > j_0$ the steady current state eventually sets on. ZRS disappears. In contrast to the above scenario, the experiments[2] demonstrate insensitive nature of the radiation-induced resistance oscillations within 30-times change in current magnitude.

Another problem in question is whether ZRS model presents a transparent physics with respect to Corbino and Hall bar simultaneous measurements[11]. For Corbino geometry sample CVC argued[10] to be N-shaped(Fig.1, top panel). At fixed voltage between the inner and outer contacts Corbino sample represents the electric field domain structure(ZCS) when $|E| < E_0$. Let us assume now ZCS at fixed applied electric field $E = E_0$ and certain value of the magnetic field. Since

$\rho_{xx} \simeq \sigma_{xx}/\sigma_{yx}^2 = 0$, the system remains ZRS as well. We argue that the experimental data[11] provide the strong evidence that the radiation-induced conductivity oscillations in Corbino samples are identical to those deduced from Hall bar measurements data. Invoking now the formalism[10], we conclude that at certain magnetic field and $E = E_0$ the ZCS(Corbino) corresponds to ZRS(Hall bar) at $j = j_0$, where $j_0 = \sigma_{yx}E_0$ according to Ref.[10]. Obviously, in real experiments[11] the above relationship is violated since the applied constant current(Hall bar setup) and constant electric field(Corbino setup) are presumably independent.

Finally, it turns out that ZRS model is unable to explain some essential experimental findings[2, 11].




FIG. 1: CVC in Corbino(top) and Hall bar(bottom) geometry suggested in Ref.[5, 10]. The bold lines depict ZCS and ZRS.

This work was supported by RFBR(grant 03-02-17588), and LSF(HPRI-CT-2001-00114, Weizmann Institute)

-
- [1] M.A.Zudov, R.R.Du, J.A.Simmons, and L.Reno, Phys. Rev. B **64**, 201311(R), 2001, M.A.Zudov, R.R.Du, L.N.Pfeiffer, and K.W.West, Phys. Rev. Lett., **90**, 046807, 2003
 - [2] R.G.Mani, J.H.Smet, K. von Klitzing, V.Narayanmurty, V.Umansky, Nature **420**, 646, 2002
 - [3] J.C.Phillips, cond-mat/0212416,0303181,0303184
 - [4] A.C.Durst, S.Sadchev, N.Read, S.M.Girvin, cond-mat/0301569
 - [5] A.V.Andreev, I.L.Aleiner, A.J.Millis, cond-mat/0302063
 - [6] P.W.Anderson, W.F.Brickman, cond-mat/0302129
 - [7] A.A.Koulakov, M.E.Raikh, cond-mat/0302465
 - [8] J.Shi, X.C.Xie, cond-mat/0302393
 - [9] S.A.Mikhailov, cond-mat/0303130
 - [10] F.S.Bergeret, B.Huckestein, A.F.Volkov, cond-mat/0303530
 - [11] C.L.Yang, M.A.Zudov, T.A.Knuuttila, R.R.Du, L.N.Pfeiffer, K.W.West, cond-mat/0303472